

# Buprevet®

## **Buprevet® - buprenorphine from Virbac**

- Potent, high-affinity, synthetic opioid agonist
- Licensed in cats and dogs for post-operative analgesia and in dogs for the potentiation of sedative effects of centrally-acting agents
- Long duration of action at 4 8 hours
- Give pre-operatively to allow analgesia at time of surgery, but also as part of a pre-emptive approach to analgesia
- Its potentiation means less sedation is needed in premedication and in anaesthesia, and recoveries can be smoother with more comfortable patients
- Versatile it is suitable for use as part of a multimodal approach to analgesia - Virbac feature Inflacam® and Carprox® in our NSAIDs range

Available in a 10 ml multidose bottle that can be broached up to 25 times and open for up to 28 days



# Introducing

# Buprevet®2ml

A breakthrough in preservative-free analgesic delivery

#### NO SNAP NO STRESS NO WASTE NO FUSS

- Buprenorphine 0.3mg/ml solution for injection in a convenient 2ml preparation containing no
- The only preservative-free buprenorphine licensed in dogs and cats for both intravenous and intramuscular administration
- Unique bunged design replaces inconvenient snap-top vials - no glass to break and a 24 hour broach life

Available in boxes of 6 x 2ml bottles.



Buprevet® 0.3 mg/ml - solution for injection. Qualitative and quantitative composition: each ml contains: active substance : buprenorphine (as hydrochloride) 0.3 mg. Excipients: glucose monohydrate, hydrochloric acid (for pH adjustment), sodium hydroxide (for pH adjustment), water for injection.

Legal Category: UK POM-V 🔊 Schedule 3

Target species: Dogs and cats.

Indications for use: Dog: Post-operative analgesia. Potentiation of the sedative effects of centrally-acting agents. Cat: Post-operative analgesia.

**Contraindications:** Do not use in case of hypersensitivity to the active substance or to any of the excipients. Do not administer by the intrathecal or peridural route. Do not use pre-operatively for caesarian section. UK: Vm 22080/4005

## BUPREVET® 10ml Multidose

Buprevet® (buprenorphine) 0.3 mg/ml solution for injection. Legal Category: UKPOM-V 🔊 Schedule 3

Target species: Dogs and cats.

Indications for use: Dog: post-operative analgesia; potentiation of the sedative effects of centrally-acting agents. Cat: post-operative analgesia.

UK: Vm 22080/4005

### TORPHADINE®

Torphadine® 10 mg/ml solution for injection. Legal Category: UK POM-V

Target species: Dogs, cats and horses.

**Indications for use:** Specifying the target species: Dog: As an analgesic: For relief of mild to moderate visceral pain. For sedation: When used in combination with certain  $\alpha 2$ -

adrenoceptor agonists (medetomidine). As a premedicant prior to general anaesthesia: For use in combination with acepromazine to provide analgesia and

sedation prior to induction of general anaesthesia. A dose-related reduction in the dose of induction-anaesthetic agent (propofol or thiopentone) is also provided.

For premedication: Give as the sole pre-anaesthetic agent. For anaesthesia: When used in combination with medetomidine

and ketamine Cat: As an analgesic for the relief of moderate pain: For pre-operative use to provide analgesia during surgery;

for post-operative analgesia after small surgical procedures. For sedation: When used in combination with certain α2-adrenoceptor agonists (medetomidine).

For anaesthesia: When used in combination with medetomidine and ketamine, suitable for short painful anaesthetic procedures. UK: Vm 41821/4037

## PROPOFOL®-LIPURO VET

Propofol®-Lipuro Vet (propofol) 10 mg/ml emulsion for injection.

Legal Category: UK POM-V Target species: dogs and cats.

**Indications for use:** Anaesthetic for dogs and cats.

UK: Vm 03551/4001

## Zoletil® 100

Zoletil® (50 mg/ml+50 mg/ml) lyophilisate and solvent for solution for injection, contains and tiletamine and zolazepam. Legal Category: UK POM-V

Target species: Dogs and cats. Indications for use: General anaesthesia.

UK: Vm 05653/4194

#### MEDETOR®

Medetor® (medetomidine hydrochloride) 1 mg/ml solution

Legal Category: UK POM-V

Target species: Dogs and cats.

Indications for use: In dogs and cats: sedation to facilitate handling; premedication prior to general anaesthesia. In cats: in combination with ketamine for general anaesthesia for minor surgical procedures of short duration.

UK: Vm 20916/4001

## REVERTOR®

Revertor® (atipamezole hydrochloride) 5 mg/ml solution for injection.

Legal Category: UK POM-V

Target species: Dogs and cats.

**Indications for use:** Atipamezole hydrochloride is a selective α2-Antagonist and indicated for reversal of the sedative effects of medetomidine and dexmedetomidine in dogs and cats.

UK: Vm 20916/4003

## **VETFLURANE®**

Vetflurane® (isoflurane) 1000 mg/g Inhalation vapour, liquid for horses, dogs, cats, ornamental birds, reptiles, rats, mice, hamsters, chinchillas, gerbils, guinea pigs and ferrets. Legal Category: UK POM-V

Indications for use: Induction and maintenance of general anaesthesia.

UK: Vm 05653/4166

See individual product SPCs for more information.

# Introducing Buprevet®2ml

## to the Virbac anaesthetics range

A breakthrough in preservative-free analgesic delivery

Propofol®-Lipuro Vet • Buprevet® • Zoletil®

Vetflurane® • Medetor® • Revertor® • Torphadine®



Use medicines responsibly. www.noah.co.uk/responsible.

Telephone: 01359 243243. Email: enquiries@virbac.co.uk www.virbac.co.uk





# Propofol<sup>®</sup>-Lipuro Vet

## Propofol®-Lipuro Vet - propofol from Virbac

- 10mg/ml emulsion for injection
- Short-acting, intravenous, induction and short-term maintenance general anaesthetic for use in dogs and cats.
  Where maintenance of anaesthesia is required, it is provided by incremental doses to effect or by inhalation anaesthetics
- Contains no preservatives so use can be repeated, even in cats; see data sheet for more information
- No age restrictions for use and used successfully for induction prior to Caesarean section

Available in boxes of 10 x 20 ml bottles

# Zoletil® 100

## Zoletil® 100 - tiletamine & zolazepan from Virbac

- 250mg tiletamine & 250mg zolazepam, powder for reconstitution
- A dissociative general anaesthetic for cats and dogs given by intramuscular or intravenous routes, Zoletil contains the active ingredients tiletamine (an NMDA receptor antagonist pharmacologically similar to ketamine) and zolazepam (a benzodiazepine) enabling complementary actions to induce the following effects:

Zoletil® 100

- Fast catalepsy followed by muscle relaxation
- Moderate superficial, immediate and visceral analgesia
- Surgical anaesthesia with muscle relaxation, preservation of laryngeal, pharyngeal and palpebral reflexes, without bulbar depression

Available as a box containing 1 x vial of lyophilisate with 1 x vial (5ml) solvent



## Vetflurane®

## **Vetflurane® - isoflurane from Virbac**

- Used for the smooth and controllable induction and maintenance of general anaesthesia
- Versatile and suitable for use in most anaesthetic protocols
- Licensed in cats and dogs (for other species see reverse)

Available in a 250 ml bottle



# Medetor® & Revertor®

## Medetor® - medetomidine from Virbac

- Potent α2 adrenergic receptor agonist licensed for intramuscular use in cats and intramuscular and intravenous use in dogs
- Medetor is used in cats and dogs for sedation, premedication to anaesthesia and with ketamine for short-acting general anaesthesia in cats
- Maximal effect within 15-20 minutes with clinical effects lasting 30-180 minutes (dose dependent)

Available in a 10 ml multi-use bottle

## Revertor® - atipamazole from Virbac

- Highly selective α2 adrenergic receptor antagonist licensed for intramuscular use in cats and dogs
- Revertor is licensed to reverse the sedative effects of medetomidine or dexmedetomidine in cats and dogs.
  It is quickly absorbed and should be given 15 to 60 minutes after medetomidine or dexmedetomidine

Available in a 10 ml multi-use bottle







## Torphadine® - butorphanol from Virbac

- 10 mg/ml solution for injection for dogs, cats and horses
- Centrally acting opioid for use as an analgesic and sedative agent for dogs and cats and as part of a multimodal approach; approved for use in combination with medetomidine, ketamine and acepromazine
- Licensed for intravenous, intramuscular and subcutaneous use in dogs and cats.

Available as a 10ml multidose bottle with a 28 day broach life





